



Evaluation of Foxtail Millet Varieties for Thermotolerance at Seedling Stage using Temperature Induction Response Technique (TIR)

Karanam Navya Jyothi ^{a++*}, V. Sumathi ^{a#},
D. Subramanyam ^{b†}, P. Sudhakar ^{a‡}
and T. Giridhar Krishna ^{a‡}

^a Regional Agricultural Research Station, Tirupati, Acharya N.G. Ranga Agricultural University, Lam, Guntur, India.

^b Department of Agronomy, S.V. Agricultural College, Tirupati, Acharya N.G. Ranga Agricultural University, Lam, Guntur, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijecc/2024/v14i114587>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126456>

Original Research Article

Received: 10/09/2024

Accepted: 13/11/2024

Published: 18/11/2024

⁺⁺ Scientist;

[#] Associate Director of Research;

[†] Professor;

[‡] Principal Scientist (Crop Physiology);

*Corresponding author: E-mail: K.navyajyothi@angrau.ac.in;

Cite as: Jyothi, Karanam Navya, V. Sumathi, D. Subramanyam, P. Sudhakar, and T. Giridhar Krishna. 2024. "Evaluation of Foxtail Millet Varieties for Thermotolerance at Seedling Stage Using Temperature Induction Response Technique (TIR)". *International Journal of Environment and Climate Change* 14 (11):785-92. <https://doi.org/10.9734/ijecc/2024/v14i114587>.

ABSTRACT

In the present scenario of climate change, Foxtail millet is the best crop of choice and an indispensable component of dryland farming due to its C₄ plant type and tolerance against the abiotic stresses. However, all the foxtail millet varieties may not sustain under severe temperature stress particularly at germination and seedling growth stages. Therefore, screening of varieties is required to identify the stress tolerant varieties in foxtail millet which can combat temperature stress at seed germination and early seedling growth stages which is the need of the hour under ever changing climatic conditions. Hence, A lab experiment was conducted to standardize the temperature induction response (TIR) protocol for foxtail millet seedlings using WGC-450 programmable plant growth chamber. Temperatures were standardized as sub lethal i.e. challenging temperatures as 38°C – 54°C for 5 hours and lethal temperatures as 56°C for 3 hours. Seven foxtail millet varieties (Suryanandi, SiA 3156, Srilakshmi, SiA 3085, Narasimharaya, Krishnadevaraya and Prasad) were screened for intrinsic tolerance using the standardized Thermo Induced Response (TIR) protocol. Out of the seven varieties evaluated, SiA 3085, Prasad, Suryanandi, Srilakshmi were identified to possess high level of thermo tolerance in terms of higher seedling survival with less reduction in root and shoot growth. The varieties with intrinsic heat tolerance can be explored for the development of heat stress tolerant varieties.

Keywords: Temperature induction response technique; heat stress; thermo tolerance; foxtail millet.

1. INTRODUCTION

“Foxtail millet is the world's second most produced millet crop and it is grown in 26 countries (Sharma and Niranjan, 2018). It plays an important role in global agriculture by feeding millions of people in arid and semi-arid locations” (Jyothi et al., 2021). “Climate change makes it the ideal crop for dryland farming due to its C₄ plant type, resistance to abiotic stresses, ability to regenerate after stress, low input requirements, ease of cultivation, and resistance to pests and diseases. However, all foxtail millet cultivars may not withstand significant moisture and temperature stress, especially during the germination and seedling growth stages. Developing screening technologies to identify thermo-tolerant foxtail millet varieties is crucial as global temperatures rise. In this context, a lab experiment was carried out to standardize the temperature induction response (TIR) procedure for foxtail millet seedlings in the WGC-450 programmable plant growth chamber. The TIR method is developed from the widely accepted principle of Lethal Dose₅₀ (LD₅₀), proposed by Trevan in 1921 for the biological standardization of insulin, toxins and drugs. Still, nowadays, it is widely used in clinical research and toxicology to determine the toxicity of drugs, pesticides and fungicides” [Pillai et al., 2021]. “The temperature induction response (TIR) technique is a robust and widely recognized empirical and non-destructive method for rapidly assessing the heat tolerance in crop plants at the seedling stage” [Raghavendra et al., 2017]. Based on this novel temperature induction response technique,

genetic variability for cellular level tolerance has been validated in many crop species by Venkatachalayya et al. (2001) in pea, Gangappa et al. (2006) in groundnut, Sudhakar et al. (2012), Harihar (2014) and Vijayalakshmi et al.(2015) in rice, Senthilkumar et al. (2004) in sunflower, Venkatesh et al. (2013) in ragi, Kheir et al. (2012) in cotton, Sairekha et al. (2016) in mung bean, Raghavendra et al. (2017) in chickpea, Ange et al. (2016) in soy bean. In the present investigation, an attempt was made to standardize the temperature induction response technique in foxtail millet which is necessary for screening the thermotolerant foxtail millet varieties which can combat temperature stress especially at seed germination and early seedling growth stages which is the need of the hour under changing climatic conditions.

2. MATERIALS AND METHODS

Research work was carried at Institute of Frontier Technology located at Regional Agricultural Research Station, Acharya N G Ranga Agricultural University, Tirupati District Andhra Pradesh with seven popular varieties of foxtail millet (SiA 3085, Prasad, Suryanandi, Krishnadevaraya, Narasimharaya SiA 3156 and Srilakshmi) which were collected from Regional Agricultural Research Station, Nandyal.

2.1 Screening of Foxtail Millet Varieties for Heat Tolerance by TIR Technique

Temperature Induction Response (TIR) technique has been developed to identify and

select the thermo tolerant genotypes. It involves exposing of seedlings or plants to induced heat stress and subsequently challenging with severe temperature and selection is based on seedling survival at the end of recovery period. This approach of TIR involves, first the identification of challenging temperature and induction temperature and later standardizing them before using for screening the material for intrinsic stress tolerance.

2.1.1 Identification of lethal temperature treatment

To assess the challenging temperatures for 100 per cent mortality, 3 days old foxtail millet seedlings of suryanandi variety were exposed to different lethal temperatures (50°C, 52°C, 54°C, 56°C and 58°C) for varying durations (1, 2 and 3 hours) without prior induction. Thus, exposed seedlings were allowed to recover at 30°C and 60 per cent relative humidity for 48 hours (Fig. 1). At the end of recovery period the temperature at which 90% mortality of the seedlings occurred was taken as the challenging temperature to assess the genetic variability for seedling survival. Percent mortality of foxtail millet genotypes after recovery was recorded. The

lethal temperature of 56°C for 3 hours was considered in this context, as maximum mortality (93%) of seedlings (Table 1).

2.1.2 Identification of sub lethal (induction) temperature

During the induction treatment, the seedlings were exposed to a gradual increase in temperature for a specific period. The temperature regimes and duration varies from crop to crop and need to be standardized. To assess and standardize the sub lethal (induction) temperature, the germinated foxtail millet seedlings (72hour old seedlings) were subjected to gradually increasing temperatures for a period of five hours. After this induction treatment, seedlings were exposed to lethal temperature i.e., 56°C for three hours and then transferred to the normal temperature for recovery. The temperature regimes and durations varied to arrive at optimum induction protocol. The optimum sub lethal temperatures were arrived based on the per cent survival of seedlings. The sub lethal treatment which recovered least per cent seedlings survival reduction was considered as optimum range of temperatures i.e., 38°C-54°C (Table 3).




Fig. 1. Protocol of the technique: Temperature Induction Response (TIR)

Table 1. Per cent mortality of foxtail millet seedlings at different lethal temperatures

S.No.	Temperature (°C)	Per cent mortality of foxtail seedlings afterrecovery		
		Duration oftemperature		
		1 hour	2 hour	3 hour
1	50	0	0	7
2	52	0	0	23
3	54	0	33	45
4	56	35	66	93
5	58	49	94	100

Table 2. Percent survival of foxtail millet seedlings different induction (sub lethal) temperature range

S. No.	Temperature range (Induction treatment for 5 h) °C	Per cent survival of seedlings
1	34-50	77
2	34-51	82
3	36-52	84
4	36-53	86
5	38-54	92
6	38-55	76

A lethal temperature of 56°C for 3 hours and induction treatment from 38-54°C for five hours was standardized using TIR (Thermo Induction Response) and considered as best lethal and induction temperatures for phenotyping of foxtail millet seedlings for intrinsic heat tolerance at cellular level.

2.2 Assessment of Thermo Induction Response (TIR)

Foxtail millet seeds were surface sterilized by treating with 0.1 per cent carbendazim for 30 minutes and washed with distilled water for 4 to 5 times followed by 1 per cent sodium hypochlorite for 1 to 2 minutes. Then the seeds were washed with distilled water for 4 to 5 times and kept for germination at 30°C and 60 per cent relative

humidity in the incubator. After 72 hours, uniform seedlings were selected in each variety and sown in aluminum trays (50 mm) filled with soil.

These trays with seedlings were subjected to sub-lethal temperatures (gradual temperature increasing from 38°C to 54°C for 5 hours in the environmental chamber (WGC-450 Programmable Plant Growth Chamber). Later these seedlings were exposed to lethal temperatures (56° C) (induced) for 3 hours. Another sub set of seedlings were directly exposed to lethal temperatures (noninduced). Induced and non-induced foxtail millet seedlings were allowed to recover at 30° C and 60 percent relative humidity for 48 hours. A control tray was maintained at 30°C, without exposing to sub-lethal and lethal temperatures.

The following parameters were recorded at 7 days after treatment from the seedlings.

a) Per cent survival of seedlings=

$$\frac{\text{No. of seedlings survived at the end of recovery}}{\text{Total number of seedlings sown in the tray}} \times 100$$

b) Per cent reduction in root growth=

$$\frac{\text{Actual root growth of control seedlings} - \text{actual root growth of treated seedlings}}{\text{Actual root growth of control seedlings}} \times 100$$

c) Per cent reduction in shoot growth=

$$\frac{\text{Actual shoot growth of control seedlings} - \text{actual shoot growth of treated seedlings}}{\text{Actual shoot growth of control seedlings}} \times 100$$

3. RESULTS

A set of seven foxtail millet varieties viz., SiA 3085, Prasad, Suryanandi, Krishnadevaraya, Narasimharaya SiA 3156 and Srilakshmi were screened for high temperature stress tolerance employing the standardized thermo induction response (TIR) protocol (Fig. 1). The experimental data was recorded and presented in Table 4 and depicted in Fig. 2.

Table 3. Performance of foxtail millet varieties for thermo induction response (TIR) characters

S.No.	Varieties	Percent survival of seedlings*	% reduction in shoot growth			% reduction in root growth		
			Actual shoot growth in control	Actual shoot growth in treatment	Percent reduction in shoot growth*	Actual root growth in control	Actual root growth in treatment	Percent reduction in root growth*
1	Suryanandi	76(93)	4.80	3.37	33.1(29.9)	4.79	3.87	23.2(19.2)
2	SIA 3156	57(70)	4.85	3.37	40.4(42.1)	4.16	2.36	41.1(43.3)
3	Srilaxmi	67(85)	3.88	2.24	33.4(30.5)	4.42	2.70	38.6(39.0)
4	SiA 3085	81(95)	4.03	3.10	27.8(23.0)	4.38	3.76	21.2(14.2)
5	Narasimharaya	60(75)	5.21	3.30	37.2(36.6)	4.88	3.47	32.5(28.9)
6	Krishnadevaraya	59(72.5)	4.10	2.55	37.9(37.8)	4.40	2.94	35.1(33.3)
7	Prasad	76(93)	4.09	3.19	30(21.9)	3.90	3.29	22.0(15.8)
	SEm ±	3.91	-	-	1.4	-	-	1.6
	CD (P=0.05)	11.7	-	-	4	-	-	4

*Data was subjected to arcsine transformation. Figures in parenthesis are original values




Fig. 2. Bar graph showing % survival of seedlings

Table 4. Identification of promising stress tolerant foxtail millet varieties through TIR technique

Tolerance Range	Varieties
Highly Stress Tolerant varieties	SiA 3085 Prasad Suryanandi Srilaxmi
Highly Stress Sensitive varieties	Krishnadevaraya Narasimharaya SiA 3156

The varieties showed significant genetic variability for per cent survival of seedlings, per cent reduction in root growth and shoot growth, respectively. The present investigation also revealed that the temperature induction response (TIR) technique can be employed in foxtail millet crop, four foxtail millet varieties viz., SiA 3085, Prasad, Suryanandi, Srilakshmi were identified to possess higher level of thermo tolerance. They showed 85-95 per cent survival of seedlings, 14.3 to 39 per cent reduction in root growth and 21.9 to 30.5 per cent reduction in shoot growth and the remaining three varieties viz., Narasimharaya, Krishnadevaraya, and SiA 3156 showed 70-85 per cent survival of seedlings, 33.3 to 43.3 per cent reduction in root growth and 36.6 to 42.1 per cent reduction in shoot growth. Thus, these varieties can be considered as highly heat sensitive varieties. Based on the observations from this technique, four varieties were classified as highly tolerant varieties and

the three varieties were classified as sensitive varieties.

4. DISCUSSION

“In spite of exposing to lethal temperature of 56°C, germination and seedling growth were less affected in the varieties viz., SiA 3085, Prasad, Suryanandi and Srilakshmi which might be probably due to acquired thermo tolerance. The results are in line with the findings of several studies, which showed that acclimated plants survive upon exposure stress, which otherwise could be lethal and are considered as heat tolerant” (Senthilkumar et al., 2004).

The remaining three varieties viz., Krishnadevaraya, Narasimharaya and SiA 3156, the seedling survival, shoot and root growth were more affected despite of the recovery conditions maintained after exposing to sub lethal to lethal

Temperature. Thus, these varieties can be considered as highly heat sensitive varieties.

5. CONCLUSION

Critical perusal of the data pertaining to the experiment revealed that the varieties SiA 3085, Prasad, Suryanandi, Srilakshmi were highly tolerant to heat stress. These varieties with intrinsic heat tolerance can be explored for the development of heat stress tolerant varieties. The results also revealed that the TIR technique is a powerful and constructive technique for identifying the foxtail millet varieties for their genetic variability in thermo tolerance within a short period of time and it is suitable for screening a large number of genotypes.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

I extend my sincere thanks to the Acharya N.G. Ranga Agricultural University, Lam, Guntur, India for providing all facilities to carry out this research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

Ange U.M., Srividhya S., Vijayalakshmi C., Boominathan P. (2016). Temperature induction response reveals intrinsic thermotolerant genotypes in soybean. *Legume Res - An Int J.* 2016;39:926-930.

Bheemesh S.J., Rao M.S., Sekhar M.R., Sudhakar P. Screening of thermotolerant foxtail millet genotypes at seedling stage using thermal induction response technique (TIR). *Agric Sci.* 2018;4(1):27-31.

Gangappa, E., Ravi, K. and Veera Kumar, G. N. (2006). Evaluation of groundnut (*Arachis hypogaea* L.) genotypes for temperature tolerance based on Temperature Induction Response (TIR) technique. *Indian Journal of Genetics and Plant Breeding.* 66(2): 127-130.

Harihar S., (2014). Temperature induction response technique – A physiological approach to identify thermotolerant genotypes in rice. *Int J Agric Sci.*; 10(1): 230-232.

Jyothi, K.N., Sumathi, V., Subramanyam, D., Sudhakar, P., Krishna, T. G., Sagar, G.K. 2021. Productivity Enhancement of Foxtail Millet (*Setaria italica* L.) through Critical Agronomic Interventions under Southern Agro-Climatic Zone of Andhra Pradesh. *International Journal of Plant & Soil Science.* 33(14): 45-51.

Kheir E.A., Sheshshayee M.S., Prasad T.G., Udayakumar M. (2012). Molecular biology and physiology: Temperature induction response as a screening technique for selecting high temperature-tolerant cotton lines. *J Cotton Sci.* 16(3):190- 199.

Pillai S., Kobayashi K., Michael M., Mathai T., Sivakumar B., Sadasivan P. (2021). John William Trevan's concept of Median Lethal Dose (LD50/LC50) – More misused than used. *J Pre-Clin Clin Res.* 15(3):137-141.

Raghavendra T., Jayalakshmi V., Babu D.V. (2017). Temperature induction response (TIR) - A novel physiological approach for thermotolerant genotypes in chickpea (*Cicer arietinum* L.). *Indian J Agric Res.*;51(3):252-256.

Sairekha, K., Reddy, D. M and Sudhakar, P. (2016). Standardization of optimum temperature conditions and screening of thermotolerant mungbean genotypes using thermo induction response (TIR) technique. *The Ecoscan.* 10(3&4): 485-489.

Senthilkumar, M., Srikanthbabu, V., Mohan, R. B., Ganeshkumar, Shivaprakash, N and Udayakumar, M. (2004). Screening of inbred lines to develop a thermotolerant sunflower hybrid using the temperature induction response (TIR) technique a novel approach by exploiting residual variability. *Journal of Experimental Botany.* 54: 2569-2578.

Sharma N, Niranjana K. (2018). Foxtail millet: Properties, processing, health benefits, and uses. *Food reviews international.* 34(4):329-363.

Sudhakar, P., Latha, P., Ramesh Babu, P., Sujatha, K and Raja Reddy, K. (2012). Identification of thermotolerant rice genotypes at seedling stage using TIR technique in pursuit of global warming. *Indian Journal of Plant Physiology.* 27: 185-188.

- Venkatachalayya., Srikanthbabu., Ganeshbabu., Krishna., P. B. T., Ramaswamy, G. K., Madappa, S and Kumar, M. K. (2001). Identification of pea genotypes with enhanced thermotolerance using temperature induction response (TIR) technique. *Journal of Plant Physiology*. 159(5): 535-545.
- Venkatesh, B. B., Sudhakar, P and Sharath, K. R. Y. (2013). Screening of thermotolerant ragi genotypes at seedling stage using TIR technique. *The Bioscan*. 8(4): 1493- 1495.
- Vijayalakshmi D., Srividhya S., Vivitha P., Raveendran M. (2015). Temperature induction response (TIR) as a rapid screening protocol to dissect the genetic variability in acquired thermotolerance in rice and to identify novel donors for high temperature stress tolerance. *Indian J Plant Physiol.*;20(4):368-374.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/126456>