

International Journal of Environment and Climate Change

Volume 14, Issue 11, Page 920-928, 2024; Article no.IJECC.127296 ISSN: 2581-8627 (Past name: British Journal of Environment & Climate Change, Past ISSN: 2231–4784)

Geospatial Analysis of LULC in the Srinagar Region of Jammu and Kashmir using Sentinel-2 Data

Khushnooda Anjum ^{a*}, Akhlaq Amin Wani ^a, M. A Islam ^a, Shah Murtaza ^a, Asif Ali Gattoo ^a, M. Iqbal Jeelani ^a and Khursheed A Sofi ^a

^a Division of Natural Resource Management, Faculty of Forestry, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Benhama Ganderbal J&K-191201, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/ijecc/2024/v14i114598

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/127296

Original Research Article

Received: 15/09/2024 Accepted: 20/11/2024 Published: 25/11/2024

ABSTRACT

Geospatial analysis based LULC mapping finds great utility in Himalyan region in order to carry proper assessment, evaluation and sustainable management of the existing resources. With the aid of remote sensing (RS) and geographic information system (GIS), the current study on LULC mapping was conducted in district Srinagar of Jammu and Kashmir for the year 2022. European Space Agency (ESA) based Sentinel-2 satellite was used as a data source. By using the technique of visual image interpretation, the study area was demarcated into 11 LULC classes viz, agriculture, forest, forest scrub, built-up, TOF, grassland, wasteland, waterbody, wetland, snow and horticulture. The findings of the study revealed that from the delineated LULC classes highest

*Corresponding author: E-mail: khushnoodaanjum345@gmail.com;

Cite as: Anjum, Khushnooda, Akhlaq Amin Wani, M. A Islam, Shah Murtaza, Asif Ali Gattoo, M. Iqbal Jeelani, and Khursheed A Sofi. 2024. "Geospatial Analysis of LULC in the Srinagar Region of Jammu and Kashmir Using Sentinel-2 Data". International Journal of Environment and Climate Change 14 (11):920-28. https://doi.org/10.9734/ijecc/2024/v14i114598.

percentage (19.61%) of area was covered by forest and least (2.17%) was covered by wetland. Moreover, the values obtained for overall classification accuracy and kappa statistics were 89.09% and 0.88 respectively. Due to increasing population and urbanisation some of the important LULC classes like wetland, agriculture etc have been encroached thereby disturbing the overall ecological balance. Therefore, findings of the current study might be useful in planning and demarcating the area where we can develop green infrastructure which might be fruitful in achieving the sustainable management goals.

Keywords: Geospatial; LULC; RS; GIS; Kashmir.

1. INTRODUCTION

Land use Land cover (LULC) are two distinct terms but are sometimes used synonymously. The social, economic and environmental factors of a place along with the human based activities over space and time play a critical role in shaping the LULC pattern of a region or place (Jamal and Ahmad. 2020). All the human related activities which take place on earth's surface is termed as Land use while as the natural or human developed features on earth's surface is referred to as Land cover (Tiwari and Kanduri, 2011). In order to put forth the management strategies to mitigate the deleterious effects of climate change, developmental activities, unprecedented and exhaustive use of natural resources on ecosystem. The periodic assessment of spatiotemporal changes in LULC classes must be incorporated as an essential tool in management plans. This will be helpful in using the various natural and man-made resources judiciously and in perpetuity along with conserving them for future generations (Maarez et al., 2022; Basheer et al., 2022). Thus, LULC mapping is essential for comprehending the different facets of the natural surroundings of humans, their activities and the sophisticated association between the two (Singh et al., 2021). It finds wide applicability in disaster management, urban and conservation planning etc (Jamal and Ahmad 2020).

Over time advancement in RS and GIS or geospatial techniques like improvement in data acquisition methods, increase in spatial, temporal and spectral resolutions have increased the utility of these techniques in analysing and monitoring LULC dynamics. Some of the fundamental areas of advancements include precise mapping, modernized and diversified information or data sources, progression in algorithms used for classification, presence of diverse open access data sources and availability of techniques for integrating data from multiple sources etc (Mashala et al., 2023)

Geospatial techniques due to their extensive pertinence are being recommended as the

efficient tools for LULC mapping. These techniques provide a great opportunity for studying the earth as an integrated system by providing a comprehensive view of the planet (Darem et al., 2023). Numerous studies have been conducted by researchers on international, national and regional level on LULC mapping, type and rates of change in LULC and its impact on social, economic and ecological aspects of the human beings and earth by obtaining data from different satellites (Ghayour et al., 2021; Bhattacharya et al., 2021; Sang et al., 2021; Mehraj et al., 2021; Dibs et al., 2020; Kaya and Görgün, 2020; Saddique et al., 2020). European Space Agency (ESA) based Sentinel-2 data has been found to be reliable and exceedingly valuable source of data for developing LULC maps due to its vast spatial resolution and temporal precision (Abbas and Jabir, 2024).

Being the summer capital of Jammu and Kashmir and trading center of the Kashmir valley, the ecological situation within Srinagar city has greatly altered (Dar, 2014). Anthropogenic activities, population growth and urbanization have immensely disrupted the fundamental ecological services. The negligence in using the natural land use capabilities and managing the same have drastically affected the ecological well-being of the city (Ahmad et al., 2024). Therefore, the resources assessment on spatiotemporal basis that could make data accessible on kind, position, geographical coverage, extent and form of alteration in each LULC class is need of the hour (Murtaza et al., 2015). Hence, the current research on LULC mapping was carried in district Srinagar of Jammu and Kashmir using Sentinel-2 as data source.

2. MATERIALS AND METHODS

2.1 Study Area

Srinagar district situated in the heart of Kashmir valley is located at an altitude of 1580 m above mean sea level and between geographical coordinates of 33°58'0"N and 34°13'0"N latitude

and 74°40'0" E and 75°9'0"E longitude (Favaz. 2023). It is bordered by district Ganderbal in north, district Pulwama in south, Budgam in south-west and Bandipora in north east. The district is surrounded by a chain of Zanskar mountains and Pir Panjal Ranges. The latitudinal variation leads to wide variation in climatic conditions due to which areas of the district at high altitude witness a typical temperate climate thereby experiencing severe cold and snowfall during winter while as places at lower altitudes experience mild summers designating tropical climate. The population density of the study area is 4407 persons per square kilometer (Kuchay et al., 2016).

The methodology adopted for conducting the research is discussed below:

Procurement of satellite data: The data was sourced from European Space Agency (ESA)

Preprocessing of satellite data: With the aid of image processing software and by using ideal band combinations False Color Composite (FCC) was made by preprocessing the acquired satellite data. The colors red, green and blue were used for developing FCC. Moreover, for the better interpretation of LULC classes in the study area various image enhancement techniques like image sharpening, changing brightness, histrogram equalisation etc were used.

Preliminary survey of the study area: For obtaining the pioneer information about study area regarding land use, topographical features etc and the number of LULC classes to be delineated, preliminary survey of the district Srinagar was carried.

LULC map generation: ArcGIS at a mapping scale of 1:30,000 (being appropriate for a satellite data with resolution of 10 m) was used for mapping purpose. The LULC classes in which satellite data was demarcated were:

1.Forest scrub 2. Forest 3. Agriculture 4. Horticulture 5. Trees outside forest 6. Wetland 7. Wasteland 8. Grassland 9. Waterbody 10. Snow 11. Built-up

Accuracy assessment of LULC map: The accuracy assessment of generated LULC map was carried by using ground truth points collected during field trips and from google earth in case of in accessible areas. In case of each LULC class 10 ground truth points were collected.

of LULC Validation map: The user's accuracy, producer's accuracy, overall accuracy and finally KAPPA (khat coefficient) were calculated for validating the map. This was done by generating the error matrix (Congalton et al., 1983).

Number of correctly classified pixels in each category $Producer's Accuracy = \frac{1}{Total number of validation points used for that category (column total)}$

Number of correctly classified pixels in each category $User's Accuracy = \frac{1}{Total number of validation points used for that category (row total)}$

 $\textbf{Overall Accuracy} = \frac{\textit{Number of correctly classified pixels}}{\textit{Total number of validation points}}$

$$k = \frac{N\sum_{i=1}^{r} x_{ij} - \sum_{i=1}^{r} (x_i + x_{ij})}{N^2 - \sum_{i=1}^{r} (x_i + x_{x+i})}$$

Where.

k = Kappa coefficient r = number of rows in the error matrix x_{ii} = the number of observations in row *i* and column *j* (on the major diagonal) x_{i} total number of observations in row *i* (shown as marginal total to right of the matrix) x_{i} total number of observations in column *i* (shown as marginal total at bottom of the matrix) N = no. of observations

Kappa statistics is a statistical tool used to reduce the error due to chance. Its value ranges from -1 to +1. It is used to measure the extent to which the data collected during an investigation or research is in agreement with the variables studied.

3. RESULTS

3.1 Land use Land Cover (LULC) of District Srinagar

The study area was delineated into 11 LULC classes among which the highest area was occupied by forest (19.61%) followed by (16.85%) by built -up, (11.33%) by agriculture, (10.04%) by forest scrub, (8.44) by wasteland, (8.25%) by grassland, (6.52%) by TOF, (6.37%) by horticulture, (6.26%) by snow, (4.16%) by waterbody and (2.17%) by wetland as shown in Table 1.The Landuse/Landcover map of district Srinagar is presented in Fig. 1. Different color code schemes were used to designate the demarcated LULC classes.

3.2 Accuracy Assessment and Validation of LULC Map

For accuracy assessment total of 110 ground truth points were collected across the study area.

The error matrix generated for the developed map of district Srinagar (2022) is shown in Table 2. This was employed for calculating producer's accuracy, user's accuracy of each LULC class. The values for overall accuracy and kappa statistics obtained were 89.09% and 0.88 respectively.

4. DISCUSSION

The current research which was undertaken in district Srinagar by using ESA based Sentinel-2 for the year 2022 as data source. For mapping purpose ArcGIS at a mapping scale of 1:30,000 was employed and the findings revealed that the maximum area was under Forest (19.61%) > Built-up (16.85%) > Agriculture (11.33%) > Forest scrub (10.04%) > Wasteland (8.44%) >

Grassland (8.25%) > TOF (6.52) >Horticulture (6.37%) > Snow (6.26) >Water body (4.16) > Wetland (2.17). Using the data from wide variety of satellites various researchers have conducted LULC assessment like the studies conducted by Raza et al. (2024), Tempa et al. (2024), Mishra et al. (2020). Fayaz, 2023 carried LULC mapping in district Srinagar by using Sentinel-2 data for the year 2020.While comparing the results with our findings it was observed that in case of LULC classes namely, built-up, forest scrub, grassland and wasteland there has been an increase of 6 ha, 2 ha, 3 ha and 2 ha respectively. However, area under LULC classes forest, snow and wetland have shown a decrease of 2 ha, 2 ha and 2.5 ha respectively. Lohare et al. (2024) carried GIS based LULC mapping in Jabalpur district, Madhva Pradesh and found that maximum area 59.26% was occupied by agriculture followed by 18.28% by forest, 18.08% hv open/barren/wasteland, 2.68% hv waterbodies and 1.70% by built-up. Hedge et al (2023) conducted an investigation on LULC practices in district Kalaburgi, Karnataka by using geospatial techniques and reported that out of the 3 delineated LULC classes, area recorded under agriculture was 83% under waterbody was 11% and 6% of area was under built-up. In a RS and GIS based research conducted on LULC dynamics in Patna, Bihar by Ahmad et al (2023) for the time period 2007-2017. It was reported that in built-up an increase of 21.86% was witnessed and in cropland and vegetation a decline of 8.95% and 8.5% respectively was recorded. Zahoor et al. (2020) conducted LULC assessment in district Ganderbal of Kashmir region using Landsat (OLI) satellite data and concluded that out of the 10 delineated LULC classes namely, grassland, snow, forest scrub, forest, agriculture, waterbody, built-up wasteland, wetland and TOF maximum area (33.96%) was

Table 1. Areal distribution of LULC classes in district Srinagar (2022)

Class	Area(ha)	%
Agriculture(a)	5420.96635	11.33
Forest(f)	9388.4899	19.61
ForestScrub(fs)	4804.6186	10.04
BuiltUp(btp)	8065.8844	16.85
Grassland(g)	3947.7492	8.25
Horticulture(h)	3047.1417	6.37
Snow(s)	2998.5004	6.26
TOF(t)	3120.7068	6.52
Waterbody(wb)	1993.15561	4.16
Wetland(wl)	1040.3288	2.17
Wasteland(wsl)	4038.81	8.44
Grand Total	47866.3517	100

Anjum et al.; Int. J. Environ. Clim. Change, vol. 14, no. 11, pp. 920-928, 2024; Article no.IJECC.127296

Class	Forest	Agriculture	Forest scrub	Buit- up	Grassland	Trees outside forest	Wasteland	Wetland	Snow	Waterbody	Horticulture	Row total
Forest	9	0	0	0	0	1	0	0	0	0	0	10
Agricuiture	0	8	0	0	0	0	0	0	0	0	2	10
Forest scrub	0	0	9	0	1	0	0	0	0	0	0	10
Built-up	0	0	0	9	0	0	0	0	1	0	0	10
Grassland	0	0	1	0	9	0	0	0	0	0	0	10
Trees outside	1	0	0	0	0	8	0	0	0	0	1	10
forest												
Wasteland	0	0	0	0	1	0	9	0	0	0	0	10
Wetland	0	0	0	0	0	0	0	10	0	0	0	10
Snow	0	0	0	0	0	0	0	0	10	0	0	10
Waterbody	0	0	0	0	0	0	0	1	0	9	0	10
Horticulture	0	1	0	0	0	1	0	0	0	0	8	10
Column Total	10	9	10	9	11	10	9	11	11	9	11	110

Table 2. Error matrix for LULC map (2022)

Overall Classification Accuracy = 89.09 % Kappa statistics = 0.88



Anjum et al.; Int. J. Environ. Clim. Change, vol. 14, no. 11, pp. 920-928, 2024; Article no. IJECC. 127296

Fig. 1. LULC map (2022) of district Srinagar

under forest while as minimum area (1.35%) was wetland. Similarly, another under study conducted on analysing resource potential and quality of soil under trees outside forests by Mehraj (2018) district Ganderbal in of Kahmir region revealed that out of the demarcated LULC classes. highest area (34.43%) was occupied by forests and lowest (1.21%) was under wetland.

In the current research, an overall classification accuracy and overall kappa statistics of 89.09 % and 0.88 respectively were obtained. During a study conducted by Seyam et al. 2023 on change in LULC in Mymensingh region of Bangladesh by using Remote sensing and GIS based techniques they documented an overall classification accuracy of 87.2% for 2002 and 89.6% for 2022 and kappa statistic value 0.84 and 0.87 for respective years. Rasool et al. (2021) conducted an investigation to detect change in LULC in South region of Kashmir from 1990 to 2017 by using various geospatial techniques. Their study concluded an overall classification accuracy of 89%, 84.56% and 92.83% respectively and the values obtained for kappa statistics were 0.89, 0.84 and 0.92. Rwanga and Ndambuki (2017) while studying LULC classification and accuracy assessment in Limpopo province in South Africa using RS and GIS based tools obtained an overall accuracy of 81.7 % and Kappa value of 0.722. Our findings are in agreement with Chakraborty *et al* (2016), Traore *et al* (2020) and Mehraj (2023).

5. CONCLUSION

By using the Geospatial analysis and Sentinel 2 satellite as data source, the current study delineated the district Srinagar into 11 LULC classes out of which highest percentage of area was under forests followed by built-up and the lowest percentage of area was under wetland. The value obtained for overall classification accuracy was 89.09 % and Kappa statistics was 0.88. The findings of the current research might helpful in identifying and evaluating be environmentally sensitive zones or LULC classes which could eventually be helpful in prioritizing these zones or classes for conservation or rehabilitation. Moreover, the LULC mapping might be helpful in understanding the consequences of various developmental and economic activities like urbanization industrialization etc on ecological stability thereby providing an insight in choosing sustainable LULC patterns for the study area.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have not been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERANCES

- Abdullah Abbas, H., & Sabah Jaber, H. (2024). Accuracy assessment of land use maps classification based on remote sensing and GIS techniques. In *BIO Web of Conferences* (Vol. 97, p. 00063). EDP Sciences.
- Ahmad, W. S., Jamal, S., Sharma, A., & Malik, I. H. (2024). Spatiotemporal analysis of urban expansion in Srinagar city, Kashmir. *Discover Cities*, 1(1), 8.
- Ahmada, M. Y., Munimb, N. H., & Sekharc, S. (2023). Analysis of changes in land use land cover and seasonal land surface temperature using remote sensing data and GIS in PMC area, Patna, Bihar. *Biospectra*, *18*(1), 1–8.
- Atay Kaya, İ., & Kut Görgün, E. (2020). Land use and land cover change monitoring in Bandırma (Turkey) using remote sensing and geographic information systems. *Environmental Monitoring and Assessment*, 192(7), 430.
- Basheer, S., Wang, X., Farooque, A. A., Nawaz, R. A., Liu, K., Adekanmbi, T., Liu, S. (2022). Comparison of land use land cover classifiers using different satellite imagery and machine learning techniques. *Remote Sensing*, 14, 4978.

DOI: doi.org/10.3390/rs14194978

Bhattacharya, R. K., Das Chatterjee, N., & Das, K. (2021). Land use and land cover change and its resultant erosion susceptible level: An appraisal using RUSLE and Logistic Regression in a tropical plateau basin of West Bengal, India. Environment, Development and Sustainability, 23(2), 1411–1446.

- Chakraborty, A., Sachdeva, K., & Joshi, P. K. (2016). Mapping long-term land use and land cover change in the central Himalayan region using a tree-based ensemble classification approach. *Applied Geography*, *74*, 136–150.
- Congalton, R. G., Oderwald, R. G., & Mead, R. A. (1983). Assessing Landsat classification accuracy using discrete multivariate analysis statistical techniques. *Photogrammetric Engineering and Remote Sensing, 49*(12), 1671–1678.
- Dar, H. (2014). Potentials and problems of adventure tourism: A study of Kashmir valley. *Abhinav International Monthly Refereed Journal of Research in Management & Technology, 3*(9), 77–84.
- Darem, A. A., Alhashmi, A. A., Almadani, A. M., Alanazi, A. K., & Sutantra, G. A. (2023). Development of a map for land use and land cover classification of the Northern Border Region using remote sensing and GIS. *The Egyptian Journal of Remote Sensing and Space Science*, 26(2), 341– 350.
- Dibs, H., Hasab, H. A., Al-Rifaie, J. K., & Al-Ansari, N. (2020). An optimal approach for land-use/land-cover mapping by integration and fusion of multispectral Landsat OLI images: Case study in Baghdad, Iraq. *Water, Air, & Soil Pollution*, 231(9), 488.
- Fayaz, S. (2023). Urban green space mapping using on screen digitization and machine learning approach in district Srinagar of J and K (M.Sc. Thesis). Sher-e-Kashmir University of Agricultural Science and Technology, Srinagar, India.
- Ghayour, L., Neshat, A., Paryani, S., Shahabi, H., Shirzadi, A., Chen, W., Al-Ansari, N., Geertsema, M., Pourmehdi Amiri, M., Gholamnia, M., & Dou, J. (2021).
 Performance evaluation of Sentinel-2 and Landsat 8 OLI data for land cover/use classification using a comparison between machine learning algorithms. *Remote Sensing*, 13(7), 1349.
- Hegde, R., Mahendra Kumar, M. B., Seema, K. V., Niranjana, K. V., & Dhanorkar, B. A. (2023). Assessment of land use/land cover change detection using geospatial techniques in Mannur-1 microwatershed of Afzalpur Taluk, Kalaburgi District, Karnataka. *Geographical Analysis*, 12(1), 34–38.

- Jamal, S., & Ahmad, W. S. (2020). Assessing land use land cover dynamics of wetland ecosystems using Landsat satellite data. *SN Applied Sciences*, 2(11), 1891.
- Kuchay, N. A., Bhat, M. S., & Shafi, N. (2016). Population growth, urban expansion and housing scenario in Srinagar City, JK, India. *Journal of Geography and Regional Planning*, 9(1), 1.
- Lohare, J., Nair, R., Pandey, S. K., Sharma, S. K., & Ramakrishnan, R. S. (2024). Assessment of classification and accuracy of land use/land cover in Jabalpur District, Madhya Pradesh, India: An analysis by remote sensing and GIS application. International Journal of Environmental and Climate Change, 14(1), 136–142.
- Maarez, H. G., Jaber, H. S., & Shareef, M. A. (2022). Utilization of Geographic Information System for hydrological analyses: A case study of Karbala province, Iraq. *Iraqi Journal of Science*, 4118–4130.
- Mashala, M. J., Dube, T., Mudereri, B. T., Ayisi, K. K., & Ramudzuli, M. R. (2023). A systematic review on advancements in remote sensing for assessing and monitoring land use and land cover impacts changes on surface water resources in semi-arid tropical environments. Remote Sensing, 15, 3926. Available:https://doi.org/10.3390/rs151639 26
- Mehraj, B. (2018). Resource mapping and soil quality assessment of trees outside forests (TOF) in Ganderbal district of Jammu and Kashmir (M. Sc. Thesis). Sher-e-Kashmir University of Agricultural Sciences and Technology, Kashmir, India.
- Mehraj, B. (2023). Site suitability modelling for white willow in district Ganderbal of Jammu and Kashmir using GIS based multicriteria approach. (Ph.D. Thesis). Sher Kashmir University of Agricultural Science and Technology, Srinagar.
- Mehraj, B., Wani, A. A., Gatoo, A. A., Saleem, I., Anjum, A., Arjumand, T., Hussain, A., Murtaza, S., Farooq, A., & Islam, M. A. (2020). Assessment of land use land cover with special reference to trees outside forests (TOF) in the central region of Kashmir Himalayas using IRS LISS-IV data. Journal of Remote Sensing and Space Science, 23(3), 122-130.
- Mishra, P. K., Rai, A., & Rai, S. C. (2020). Land use and land cover change detection using

geospatial techniques in the Sikkim Himalaya, India. *The Egyptian Journal of Remote Sensing and Space Science*, *23*(2), 133–143.

- Murtaza, K. O., Rather, M. A., Lateef, Z., Shah, V., Kuchhay, N. (2015). Geospatial mapping of Srinagar City. *Journal of Geography and Regional Planning*, 8(6), 157–165.
- Rasool, R., Fayaz, A., ul Shafiq, M., Singh, H., & Ahmed, P. (2021). Land use land cover change in Kashmir Himalaya: Linking remote sensing with an indicator-based DPSIR approach. *Ecological Indicators*, 125, 107447.
- Raza, A., Syed, N. R., Fahmeed, R., Acharki, S., Aljohani, T. H., Hussain, S., Zubair, M., Zahra, S. M., Islam, A. R., Almohamad, H., & Abdo, H. G. (2024). Investigation of changes in land use/land cover using principal component analysis and supervised classification from operational land imager satellite data: A case study of underdeveloped regions, Pakistan. Discover Sustainability, 5(1), 73.
- Rwanga, S. S., & Ndambuki, J. M. (2017). Accuracy assessment of land use/land cover classification using remote sensing and GIS. *International Journal of Geosciences*, 8(4), 611– 618.
- Saddique, N., Mahmood, T., & Bernhofer, C. (2020). Quantifying the impacts of land use/land cover change on the water balance in the afforested River Basin, Pakistan. *Environmental Earth Sciences*, 79(19), 448.
- Sang, X., Guo, Q., Wu, X., Xie, T., He, C., Zang, J., Qiao, Y., Wu, H., & Li, Y. (2021). The effect of DEM on the land use/cover classification accuracy of Landsat OLI images. *Journal of the Indian Society of Remote Sensing*, *49*, 1507– 1518.
- Seyam, M. M., Haque, M. R., & Rahman, M. M. (2023). Identifying the land use land cover (LULC) changes using remote sensing and GIS approach: A case study at Bhaluka in Mymensingh, Bangladesh. *Case Studies in Chemical and Environmental Engineering*, 7, 100293.
- Singh, R. K., Sinha, V. S., Joshi, P. K., & Kumar, M. (2021). A multinomial logistic modelbased land use and land cover classification for the South Asian Association for Regional Cooperation

nations using Moderate Resolution Imaging Spectroradiometer product. *Environment, Development and Sustainability*, 23(4), 6106–6127.

- Tempa, K., Ilunga, M., Agarwal, A., & Tashi. (2024). Utilizing Sentinel-2 satellite imagery for LULC and NDVI change dynamics for Gelephu, Bhutan. *Applied Sciences*, *14*(4), 1578.
- Tiwari, K., & Kanduri, K. (2011). Land use land cover change detection in Doon Valley. Dehradun Tehsil, Uttarakhand using GIS and remote sensina technique. International Journal of Geomatics and Geosciences, 2, 11 -23.
- Traore, M., Ndepete, C. P., Zaguy-Guerembo, R. L., & Pour, A. B. (2020). Assessment of land use/land cover change mapping in Bangui city using remote sensing and GIS techniques. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 43, 1651– 1656.
- Zahoor, S., Wani, A. A., Masoodi, T. H., Islam, M. A., Gatoo, A. A., & Gangoo, S. A. (2020). Land use land cover classification and mapping using geospatial techniques in Ganderbal District, J & K, India. *International Journal of Current Microbiology and Applied Sciences*, *9*(11), 3838–3847.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/127296